www.elsevier.com/locate/bres

Author index

Allen, S.J., see Smith, S.K.F. (78) 100
Anderson, H.A., see Smith, S.K.F. (78) 100
Aruga, J., Shimoda, K. and Mikoshiba, K.
A 5' segment of the mouse *Zic1*gene contains a region specific
enhancer for dorsal hindbrain and
spinal cord (78) 15

Baas, D., Bumsted, K.M., Martinez, J.A., Vaccarino, F.M., Wikler, K.C. and Barnstable, C.J.

The subcellular localization of OTX2 is cell-type specific and developmentally regulated in the mouse retina (78) 26

Barnstable, C.J., see Baas, D. (78) 26

Bevilaqua, L., see Paratcha, G. (78) 120

Beyer, C., see Caba, M. (78) 69

Bickford, P., see Eckles-Smith, K. (78) 154

Blume, A., see Stroth, U. (78) 175

154 Bumsted, K.M., see Baas, D. (78) 26

Caba, M., Pau, K.-Y.F., Beyer, C., Gonzalez, A., Silver, R. and Spies, H.G. Coitus-induced activation of c-fos and gonadotropin-releasing hormone in hypothalamic neurons in female rabbits (78) 69

Browning, M.D., see Eckles-Smith, K. (78)

Cammarota, M., see Paratcha, G. (78) 120
Castelli, M.P., Mocci, I., Langlois, X.,
Gommeren, W., Luyten, W.H.M.L.,
Leysen, J.E. and Gessa, G.L.
Quantitative autoradiographic
distribution of γ-hydroxybutyric acid
binding sites in human and monkey
brain (78) 91

Chang, W.-T., see Wang, H.-L. (78) 146
Chen, L., Cooper, N.G.F. and Mower, G.D.
Developmental changes in the
expression of NMDA receptor
subunits (NR1, NR2A, NR2B) in the
cat visual cortex and the effects of
dark rearing (78) 196

Chen, M.-S., see Wang, H.-L. (78) 146 Chiba, T., see Ino, H. (78) 163 Choi, E.J., see Ha, C.M. (78) 59 Choi, W.S., see Ha, C.M. (78) 59 Christjansen, K.N., see Poulsen, C.F. (78) 201 Chun, S.Y., see Ha, C.M. (78) 59 Clayton, D., see Eckles-Smith, K. (78) 154 Cooper, N.G.F., see Chen, L. (78) 196 Dawbarn, D., see Smith, S.K.F. (78) 100 de Iraldi, A.P., see Paratcha, G. (78) 120 de Stein, M.L., see Paratcha, G. (78) 120 Dorsa, D.M., see McMillan, P.J. (78) 138

Ebihara, S., see Yoshimura, T. (78) 207
Eckles-Smith, K., Clayton, D., Bickford, P. and Browning, M.D.
Caloric restriction prevents age-related deficits in LTP and in NMDA receptor expression (78) 154

Fang, W., see Wang, H.-L. (78) 146 Fraser, P.E., see Smith, S.K.F. (78) 100 Fukushima, N., see Hamabe, W. (78) 186

Gessa, G.L., see Castelli, M.P. (78) 91 Goksøyr, A., see Øyan, A.M. (78) 38 Gommeren, W., see Castelli, M.P. (78) 91 Gonzalez, A., see Caba, M. (78) 69 Granier, C., see Torreilles, F. (78) 181

Ha, C.M., Kang, J.H., Choi, E.J., Kim, M.S., Park, J.-W., Kim, Y., Choi, W.S., Chun, S.Y., Kwon, H.B. and Lee, B.J. Progesterone increases mRNA levels of pituitary adenylate cyclase-activating polypeptide (PACAP) and type I PACAP receptor (PAC₁) in the rat hypothalamus (78) 59

Hamabe, W., Fukushima, N., Yoshida, A. and Ueda, H.
Serum-free induced neuronal apoptosis-like cell death is independent of caspase activity (78)

Hara, Y., see Yamamoto, Y. (78) 108 Hartvig, L., see Poulsen, C.F. (78) 201 Hastrup, S., see Poulsen, C.F. (78) 201 Ho, I.K., see Jang, C.-G. (78) 204 Holmqvist, B., see Øyan, A.M. (78) 38

Ino, H. and Chiba, T.

Expression of proliferating cell
nuclear antigen (PCNA) in the adult
and developing mouse nervous system
(78) 163

Izquierdo, I., see Paratcha, G. (78) 120

Jang, C.-G., Park, Y., Tanaka, S., Ma, T., Loh, H.H. and Ho, I.K. Involvement of μ-opioid receptors in potentiation of apomorphine-induced climbing behavior by morphine: studies using μ -opioid receptor gene knockout mice (78) 204

Kang, J.H., see Ha, C.M. (78) 59
Kato, A., see Taniguchi, M. (78) 131
Kim, M.S., see Ha, C.M. (78) 59
Kim, Y., see Ha, C.M. (78) 59
Kobayashi, A., see Taniguchi, M. (78) 131
Kobayashi, S., see Murata, S. (78) 80
Kohmura, E., see Taniguchi, M. (78) 131
Kumura, E., see Taniguchi, M. (78) 131
Kuroiwa, A., see Yoshimura, T. (78) 207
Kwon, H.B., see Ha, C.M. (78) 59

Langlois, X., see Castelli, M.P. (78) 91 Lee, B.J., see Ha, C.M. (78) 59 Leverenz, J.B., see McMillan, P.J. (78) 138 Levesque, M.F., see Palm, K. (78) 192 Leysen, J.E., see Castelli, M.P. (78) 91 Li, A.H., see Wang, H.-L. (78) 146 Loh, H.H., see Jang, C.-G. (78) 204 Lopez, M., see Paratcha, G. (78) 120 Luyten, W.H.M.L., see Castelli, M.P. (78) 91

Ma, T., see Jang, C.-G. (78) 204
Mack, K.J., see Nanda, S.A. (78) 1
Makino, E., see Yoshimura, T. (78) 207
Martinez, J.A., see Baas, D. (78) 26
Maruno, M., see Taniguchi, M. (78) 131
Mason, G., see Smith, S.K.F. (78) 100
Matsuda, Y., see Yoshimura, T. (78) 207
Matute, C., see Morán-Jiménez, M.-J. (78) 50
McMillan, P.J., Leverenz, J.B. and Dorsa, D.M.

Specific downregulation of presenilin 2 gene expression is prominent during early stages of sporadic late-onset Alzheimer's disease (78) 138

Medina, J.H., see Paratcha, G. (78) 120
Mielke, K., see Stroth, U. (78) 175
Mikoshiba, K., see Aruga, J. (78) 15
Mocci, I., see Castelli, M.P. (78) 91
Morán-Jiménez, M.-J. and Matute, C.
Immunohistochemical localization of the P2Y₁ purinergic receptor in neurons and glial cells of the central

nervous system (78) 50

Mourton-Gilles, C., see Torreilles, F. (78) 181 Mower, G.D., see Chen, L. (78) 196 Murata, S., Usuda, N., Okano, A., Kobayashi, S. and Suzuki, T.

Occurrence of a transcription factor,

signal transducer and activators of transcription 3 (Stat3), in the postsynaptic density of the rat brain (78) 80

Namikawa, T., see Yoshimura, T. (78) 207 Nanda, S.A. and Mack, K.J.

> Seizures and sensory stimulation result in different patterns of brain derived neurotrophic factor protein expression in the barrel cortex and hippocampus (78) 1

Naylor, R.L., see Smith, S.K.F. (78) 100 Neuman, T., see Palm, K. (78) 192 Nilsen, F., see Øyan, A.M. (78) 38 Nomura, M., see Yamamoto, Y. (78) 108

Ohnishi, T., see Taniguchi, M. (78) 131
Okano, A., see Murata, S. (78) 80
Øyan, A.M., Nilsen, F., Goksøyr, A. and
Holmqvist, B.
Partial cloning of constitutive and
inducible nitric oxide synthases and
detailed neuronal expression of NOS
mRNA in the cerebellum and optic
tectum of adult Atlantic salmon
(Salmo salar) (78) 38

Palm, K., Salin-Nordström, T., Levesque, M.F. and Neuman, T.
Fetal and adult human CNS stem cells have similar molecular characteristics and developmental potential (78) 192

Paratcha, G., de Stein, M.L., Szapiro, G., Lopez, M., Bevilaqua, L., Cammarota, M., de Iraldi, A.P., Izquierdo, I. and Medina, J.H.

Experience-dependent decrease in synaptically localized Fra-1 (78) 120

Park, J.-W., see Ha, C.M. (78) 59

Park, Y., see Jang, C.-G. (78) 204 Pau, B., see Torreilles, F. (78) 181 Pau, K.-Y.F., see Caba, M. (78) 69 Poulsen, C.F., Christjansen, K.N., Hastrup, S. and Hartvig, L. Identification and cloning of a γ3 subunit splice variant of the human GABA_A receptor (78) 201

Robertson, A.G.S., see Smith, S.K.F. (78) 100 Roche, P.A., see Smith, S.K.F. (78) 100 Roquet, F., see Torreilles, F. (78) 181

Salin-Nordström, T., see Palm, K. (78) 192
Serino, R., see Yamamoto, Y. (78) 108
Shibuya, I., see Yamamoto, Y. (78) 108
Shimoda, K., see Aruga, J. (78) 15
Shirahata, A., see Yamamoto, Y. (78) 108
Silver, R., see Caba, M. (78) 69
Smith, S.K.F., Anderson, H.A., Yu, G.,
Robertson, A.G.S., Allen, S.J., Tyler,
S.J., Naylor, R.L., Mason, G., Wilcock,
G.W., Roche, P.A., Fraser, P.E. and
Dawbarn, D.
Identification of syntaxin 1A as a
novel binding protein for presenilin-1
(78) 100

Spies, H.G., see Caba, M. (78) 69 Stroth, U., Blume, A., Mielke, K. and Unger, T.

Angiotensin AT₂ receptor stimulates ERK1 and ERK2 in quiescent but inhibits ERK in NGF-stimulated PC12W cells (78) 175

Suzuki, T., see Murata, S. (78) 80 Suzuki, T., see Yoshimura, T. (78) 207 Suzuki, Y., see Yoshimura, T. (78) 207 Szapiro, G., see Paratcha, G. (78) 120

Tamatani, M., see Taniguchi, M. (78) 131
Tanaka, S., see Jang, C.-G. (78) 204
Taniguchi, M., Yamashita, T., Kumura, E.,
Tamatani, M., Kobayashi, A., Yokawa,
T., Maruno, M., Kato, A., Ohnishi, T.,
Kohmura, E., Tohyama, M. and
Yoshimine, T.
Induction of aquaporin-4 water

channel mRNA after focal cerebral ischemia in rat (78) 131

Tohyama, M., see Taniguchi, M. (78) 131

Torreilles, F., Roquet, F., Granier, C., Pau, B. and Mourton-Gilles, C.

Binding specificity of monoclonal antibody AD2: influence of the phosphorylation state of tau (78) 181

Tyler, S.J., see Smith, S.K.F. (78) 100

Ueda, H., see Hamabe, W. (78) 186 Ueta, Y., see Yamamoto, Y. (78) 108 Unger, T., see Stroth, U. (78) 175 Usuda, N., see Murata, S. (78) 80

Vaccarino, F.M., see Baas, D. (78) 26

Wang, H.-L., Wu, T., Chang, W.-T., Li, A.H., Chen, M.-S., Wu, C.-Y. and Fang, W. Point mutation associated with Xlinked dominant Charcot-Marie-Tooth disease impairs the P2 promoter activity of human connexin-32 gene (78) 146

Wikler, K.C., see Baas, D. (78) 26 Wilcock, G.W., see Smith, S.K.F. (78) 100 Wu, C.-Y., see Wang, H.-L. (78) 146 Wu, T., see Wang, H.-L. (78) 146

Yamamoto, Y., Ueta, Y., Hara, Y., Serino, R., Nomura, M., Shibuya, I., Shirahata, A. and Yamashita, H. Postnatal development of orexin/ hypocretin in rats (78) 108 Yamashita, H., see Yamamoto, Y. (78) 108 Yamashita, T., see Taniguchi, M. (78) 131 Yokawa, T., see Taniguchi, M. (78) 131 Yoshida, A., see Hamabe, W. (78) 186 Yoshimine, T., see Taniguchi, M. (78) 131 Yoshimura, T., Suzuki, Y., Makino, E., Suzuki, T., Kuroiwa, A., Matsuda, Y., Namikawa, T. and Ebihara, S. Molecular analysis of avian circadian clock genes (78) 207 Yu, G., see Smith, S.K.F. (78) 100